

# NATURAL FOREST CONVERSION AND ITS IMPACT ON POPULATIONS OF KEY LIVELIHOOD TREE SPECIES IN OMO BIOSPHERE RESERVE, NIGERIA

\*Chima, U.D.<sup>1</sup> and Ihuma, J.O.<sup>2</sup>

1. Department of Forestry and Wildlife Management, University of Port Harcourt, P.M.B. 5323, Port Harcourt, Rivers State, Nigeria.

\*Corresponding author's email: [uzoma.chima@uniport.edu.ng](mailto:uzoma.chima@uniport.edu.ng)

2. Department of Biological Sciences, Bingham University, P.M.B. 005, Karu, Nasarawa State, Nigeria

## ABSTRACT

*Natural forest conversion and unsustainable use of forest resources are on the increase without adequate consideration of their implications for sustainable livelihoods. This study examined the impact of natural forest conversion on key livelihood tree species in Omo Biosphere Reserve, by examining their populations in the Strict Nature Reserve (SNR), Nauclea diderrichii Plantation (NDP), Tectona grandis Plantation (TGP), Pinus caribaea Plantation, Gmelina arborea Plantation, Theobroma cacao Plantation (CP), and three age-sequences of arable farmland – AF<sub>1</sub>, AF<sub>2</sub>, and AF<sub>3</sub>. The SNR was the most species rich ( $n = 17$ ) and diverse ( $H = 2.6210$ ; Simpson  $1-D = 0.9127$ ) of all the land use types. Key livelihood tree species diversity was higher in the arable farmlands ( $H = 0.7608$  to  $1.3810$ ; Simpson  $1-D = 0.3765$  to  $0.7111$ ) than in the monoculture plantations ( $H = 0.0313$  to  $1.311$ ; Simpson  $1-D = 0.0099$  to  $0.6701$ ) with GAP being the least diverse. The NDP was more similar to the SNR ( $SI = 21.74$ ) than any other land use type. The NDP showed a closer association with AF<sub>1</sub> and AF<sub>2</sub> in its key livelihood tree species than with other monoculture plantations. The CP was ecologically the farthest from the other land use types with respect to key livelihood tree species composition. The study showed that natural forest conversion to monoculture plantations and arable farm reduce key livelihood tree species richness and diversity, and that higher degree of disturbance as a result of high impact logging and longer period of cultivation, beyond thirty years, exacerbates the problem.*

**Key words:** Land use, deforestation, livelihood, tree diversity, monoculture, farming

## INTRODUCTION

The burgeoning population of humans in Nigeria and other developing countries has led to indiscriminate use of land resources not minding the short and long term socio-economic and ecological consequences. Deforestation has continued unabated despite

the apparently enormous environmental consequences associated with it.

Forests provide sources of livelihood like food, shelter, clothing and heating and a great majority of people living in poverty depend on forests and trees outside forests to generate income through employment and through the

sale of forest goods and services. It has been observed that more than 25% of the world's population – an estimated 1.6 billion people – rely on forest resources for their livelihoods, and of these, almost 1.2 billion live in extreme poverty (World Bank (2001); and lack the basic necessities to maintain a decent standard of living, for instance, sufficient and nutritious food, adequate shelter, access to health services, energy sources, safe drinking water, education and a healthy environment (FAO, 2006).

In Nigeria, forests provide invaluable services to the nation. But over the last half century, the Nigerian rainforest has experienced unprecedented reduction due to deforestation and degradation, which now pose intractable ecological, land use, biodiversity and sustainable management problems (Ikhuoria *et al.*, 2006). This has negative implications on rural livelihoods due to the near-absolute dependence of the rural populace on biodiversity, for their sustenance.

Chima *et al.* (2012) had documented and prioritized the key livelihood tree species in the reserve using the user preference approach. The human populations in Omo

Biosphere Reserve which is mainly rural depend to a large extent, on forest resources, for their living. However, despite the high spate of deforestation and the conversion of the natural forests to other land uses like monoculture plantations of exotic tree species, cocoa plantations and arable farms, no empirical study had been carried out to ascertain the impact on the populations of trees that support rural livelihoods.

This study therefore, examined the impact of natural forest conversion on the populations of key livelihood tree species by comparing them between a natural forest and introduced land use types in the reserve. It is hoped that the information provided in this study will enable management decisions that will enhance the conservation of the key livelihood tree species.

## **MATERIALS AND METHODS**

### **The Study Area**

Omo Biosphere Reserve is located between latitudes 6° 35' to 7° 05' N and longitudes 4° 19' to 4° 40' E in the South-west of Nigeria, and covers an area of about 130,500 hectares (Ojo, 2004). The reserve is in the mixed moist

semi-evergreen rainforest zone (Ola-Adams, 1999). However, anthropogenic activities, mainly logging, establishment of monoculture plantations, and farming, have changed the original vegetation of the reserve to a large extent. Geologically, the reserve lies on crystalline rocks of the undifferentiated basement complex which in the southern parts is overlain by Eocene deposits of sand, clay and gravel (Isichei, 1995). It has an undulating terrain with maximum elevation of 150 m

above sea level towards the west while the lowest parts of the reserve are in the south. The Lagos-Ore-Benin Highway passes through the southern tip of the reserve. The reserve falls within the tropical wet-and-dry climate characterized by two rainfall peaks separated by a relatively less humid period usually in the month of August (Ola-Adams, 1999). Figure 1 is the map of Omo Biosphere Reserve showing the study sites and surrounding reserves.




Figure 1: Map showing Omo Biosphere Reserve, the study sites and surrounding reserves  
Source: Adapted from Ola-Adams (1999)

## Selection of Study Sites

Nine sites representing different land use/land cover types were purposively chosen for the study. The histories of the sites were obtained from the Ogun State Forestry Department's Office at Area J4 of the reserve. The reference site (6.96598°N and 4.36245°E) was taken from the Strict Nature Reserve at Etemi. This site represents part of the reserve that has not been modified either by agricultural activities of the smallholders, plantation establishment or timber exploitation. Three other sites - AF<sub>1</sub> (6°50'26.77"N and 4°21'37.03"E), AF<sub>2</sub> (6°50'29.71"N and 4°21'37.61"E) and AF<sub>3</sub> (6°50'32.80"N and 4°21'38.85"E); were selected from around Mile 1 enclave in Area J4, to reflect three chronosequences of arable farmland. Sites AF<sub>1</sub>, AF<sub>2</sub> and AF<sub>3</sub> were originally established as taungya farms and have been under cultivation since they were given out to farmers in 2000, 1990, and 1975 respectively. Site CP (6°52'49.82"N and 4°24'48.91"E) was chosen from a pure Cocoa Plantation established in the year 2000, near Temidire Camp. Four other sites – *Pinus caribaea* Plantation (PCP - 6°50'03.54"N and 4°22'00.65"E); *Tectona grandis* Plantation (TGP - 6°50'08.37"N and 4°21'39.92"E); *Gmelina*

*arborea* Plantation (GAP - 6°54'13.94"N and 4°22'30.44"E); and *Nauclea diderichii* Plantation (NDP - 6°50'16.11"N and 4°22'05.56"E); were chosen to represent monoculture plantations of different species and ages. PCP was established in 1997, TGP in 1989, GAP in 1983, and NDP in 1975. PCP has not been logged since establishment but bears a pineapple orchard. TGP had been logged and now bears mainly coppices on the felled stumps. GAP had been logged extensively, though mature trees and saplings abound. NDP has not been logged since it was established.

## Data Collection

Ten 35 m × 35 m quadrats were randomly distributed in each of the sites for the enumeration of the key livelihood tree species (Table 1). This quadrat size falls within the range specified in the literature for ecological studies in the humid tropics (Salami, 2006). Narrow cut lines were used to demarcate plot boundaries. Species identification was done by an expert taxonomist from the Forestry Research Institute of Nigeria (FRIN), Ibadan, with the aid of keys provided by Keay (1989). All single-stem woody plants of erect posture with a minimum height of 5 m and diameter at breast height (dbh) of 5 cm were

identified to species level and the number of individuals counted and recorded. This tree size was considered to ensure that only mature trees were captured. Specimens of species that could

not be identified in the field were taken to the Forest Research Institute of Nigeria Herbarium, for identification.

**Table 1: Checklist of key livelihood tree species and their ranking**

S/No.	Species	Common or Local name	Family	Total Score	Rank
1	<i>Khaya ivorensis</i>	Lagos mahogany	Meliaceae	1295	1 <sup>st</sup>
2	<i>Nauclea diderrichii</i>	Opepe	Rubiaceae	1240	2 <sup>nd</sup>
3	<i>Terminalia ivorensis</i>	Black afara	Combretaceae	850	3 <sup>rd</sup>
4	<i>Cordia millenii</i>	Omo	Boraginaceae	690	4 <sup>th</sup>
5	<i>Alstonia boonei</i>	Pattern wood	Apocynaceae	465	5 <sup>th</sup>
6	<i>Terminalia superba</i>	White afara	Combretaceae	375	6 <sup>th</sup>
7	<i>Erythroleum suaveolens</i>	Erun-obo	Leguminosae - Caesalpinioideae	330	7 <sup>th</sup>
8	<i>Mangifera indica</i>	Mango	Anacardiaceae	265	8 <sup>th</sup>
9	<i>Entandrophragma utile</i>	Jebo	Meliaceae	260	9 <sup>th</sup>
10	<i>Anacardium occidentale</i>	Cashew	Anacardiaceae	260	9 <sup>th</sup>
11	<i>Milicia excelsa</i>	Iroko	Moraceae	255	11 <sup>th</sup>
12	<i>Lophira alata</i>	Ekki	Ochnaceae	190	12 <sup>th</sup>
13	<i>Triplochiton schleroxylon</i>	Obeche	Sterculiaceae	190	12 <sup>th</sup>
14	<i>Piptadeniastrum africanum</i>	Agboyin	Leguminosae - Mimosoideae	175	14 <sup>th</sup>
15	<i>Theobroma cacao</i>	Cocoa	Malvaceae	145	15 <sup>th</sup>
16	<i>Mitragyna ciliata</i>	African linden	Rubiaceae	140	16 <sup>th</sup>
17	<i>Mansonia altissima</i>	Mansonia	Sterculiaceae	140	16 <sup>th</sup>
18	<i>Ceiba pentandra</i>	Kapok tree	Malvaceae	130	18 <sup>th</sup>
19	<i>Enantia chlorantha</i>	Osopupa, Yaru	Annonaceae	130	18 <sup>th</sup>
20	<i>Cedrela odorata</i>	Honduras cedar	Meliaceae	110	20 <sup>th</sup>
21	<i>Anthonotha macrophylla</i>	Abara	Leguminosae - Caesalpinioideae	110	20 <sup>th</sup>
22	<i>Elaeis guineensis</i>	Palm tree	Arecaceae	110	20 <sup>th</sup>
23	<i>Citrus sinensis</i>	Sweet orange	Rutaceae	100	23 <sup>rd</sup>
24	<i>Cola nitida</i>	Kola nut	Sterculiaceae	90	24 <sup>th</sup>
25	<i>Buchholzia coriacea</i>	Wonderful kola	Capparidaceae	85	25 <sup>th</sup>
26	<i>Gmelina arborea</i>	Gmelina	Verbenaceae	80	26 <sup>th</sup>
27	<i>Entandrophragma angolense</i>	Ijebo	Meliaceae	75	27 <sup>th</sup>
28	<i>Nesogordonia papaverifera</i>	Danta	Sterculiaceae	55	28 <sup>th</sup>
29	<i>Newbouldia laevis</i>	Boundary tree	Bignoniaceae	55	28 <sup>th</sup>
30	<i>Citrus aurantifolia</i>	Lime	Rutaceae	55	28 <sup>th</sup>
31	<i>Garcinia kola</i>	Bitter kola	Guttiferae	40	31 <sup>st</sup>
32	<i>Azadirachta indica</i>	Neem	Meliaceae	40	31 <sup>st</sup>
33	<i>Daniellia ogea</i>	Ogea	Leguminosae - Caesalpinioideae	35	33 <sup>rd</sup>
34	<i>Tectona grandis</i>	Teak	Verbenaceae	25	34 <sup>th</sup>
35	<i>Cleistopholis patens</i>	Apako	Annonaceae	25	34 <sup>th</sup>
36	<i>Terminalia catappa</i>	Indian almond	Combretaceae	20	36 <sup>th</sup>
37	<i>Chrysophyllum albidum</i>	African star apple	Sapotaceae	15	37 <sup>th</sup>
38	<i>Parinari sp.</i>	Abere	Chrysobalanaceae	15	37 <sup>th</sup>

Source: Adapted from Chima *et al.* (2012).

## Data analysis

### Measurement of Alpha Diversity

In this study, Simpson Index (Simpson, 1949) and Shannon-Wiener Index (Odum, 1971) were used to measure the diversity of key livelihood tree species in each land use type. These indices were chosen because they provide measures of the different components of diversity. The Shannon-Wiener index reflects the manner in which abundance is distributed amongst the different species constituting the community. The index is based on the relative frequencies of species in the population (Giramet-Carpentier *et al.*, 1998), thus taking into account both species richness and evenness. However, Magurran, (1988) notes that the value of the index is most strongly related to species richness. Simpson's index is a dominance measure since it is weighted towards the abundance of the most common species in a sample rather than providing a measure of species richness. According to Magurran (1988), it reflects the probability of any two individuals drawn at random from an infinitely large population belonging to different species, and the index is less sensitive to species richness.

- Simpson's Index is expressed as:

$$D = \frac{\sum_{i=1}^q ni(ni-1)}{N(N-1)} \text{----- Eqn. 1}$$

#### Where:

N = total number of individuals encountered

ni = number of individuals of ith species

enumerated for i=1.....q

q = number of different species enumerated.

Since Simpson's index as expressed above is not directly related to diversity (i.e. the lower the index, the higher the diversity and *vice versa*), it is expressed in this study as (1 – D) to allow for a direct relationship.

- Shannon-Wiener Index is expressed as:

$$H = - \sum_{i=1}^s pi \log pi \text{----- Eqn. 2}$$

#### Where:

pi = the proportion of individuals in the ith species

s = the total number of species

Both Simpson and Shannon-Wiener diversity indices were computed using the PAleontological Chima and Ihuma Software.

### Measurement of Beta Diversity/Similarity

Beta diversity is a measure of the extent to which the diversity of two or more spatial units differs (Magurran, 2004) and is generally used to characterise the degree of spatial heterogeneity in

diversity at the landscape scale, or to measure the change in diversity along transects of environmental gradients. Wolda (1983) suggested the use of similarity indices for measuring beta diversity. However, Jansen and Vegelius (1981) observed that, of the many similarity indices, only three of them (the Ochiai, the Jaccard and the Sorensen) are worth considering. Hence, Sorensen's similarity index (Pielou, 1969) was used to determine the similarity in species composition of land use types considered in this study. Recent studies (e.g. Ogunleye *et al.*, 2004; Ojo, 2004; Ihuma *et al.*, 2011; Chima *et al.*, 2011) have also employed the Sorensen's index to measure beta diversity.

Sorensen's Similarity Index is expressed as:

$$SI = \frac{a}{a + b + c} * 100 \quad \text{----- Eqn. 3}$$

Where: a = number of species present in both land use types

b = number of species present in land use type 1 but absent in land use type 2

c = number of species present in land use type 2 but absent in land use type 1

## Cluster Analysis

Cluster analysis was performed using the PAleontological STatistics (PAST) software to provide a hierarchical classification of the various land use types, such that land use types with more similar key livelihood tree species are grouped into the same cluster while dissimilar ones are grouped into different clusters. In performing the cluster analysis, the Sorensen's similarity index was used to measure the ecological distances between land use types.

## RESULTS

### Diversity of key Livelihood Tree Species at different Land use Types

Key livelihood tree species diversity indices for all land use types are presented in Table 2. The SNR was the most diverse of all the land use types. Key livelihood tree species diversity was higher in the arable farmlands than in the monoculture plantations with GAP being the least diverse.

**Table 2: Diversity indices for key livelihood tree species in different land use types**

	SNR	GAP	CP	PCP	NDP	TGP	AF <sub>1</sub>	AF <sub>2</sub>	AF <sub>3</sub>
No. of species	17	2	4	5	12	5	7	6	4
Individuals	65	1007	1278	24	1278	1089	75	33	18
Dominance	0.0873	0.9901	0.9225	0.3299	0.8284	0.8893	0.2889	0.3939	0.6235
Shannon H	2.6210	0.0313	0.2041	1.311	0.4690	0.2683	1.3810	1.1420	0.7608
Simpson 1-D	0.9127	0.0099	0.0775	0.6701	0.1716	0.1107	0.7111	0.6061	0.3765

Source: Field Survey, 2012

### Similarity of land use types in terms of key livelihood tree species composition

Similarity and associations between land use types are shown in Table 3 and Figure 2 respectively. The NDP was more similar to the SNR than both the other monoculture

plantations and arable farmlands. The NDP showed a closer association to AF<sub>1</sub> and AF<sub>2</sub> (Figure 2). The CP was ecologically the farthest from the other land use types with respect to the key livelihood tree species.

**Table 3: Sorensen's Similarity Indices for key livelihood tree species at different sites**

	SNR	GAP	CP	PCP	NDP	TGP	AF <sub>1</sub>	AF <sub>2</sub>	AF <sub>3</sub>
SNR	*	5.56	5.00	4.76	21.74	4.76	20.00	21.00	16.67
GAP		*	0.00	16.67	7.69	16.67	28.57	33.33	0.00
CP			*	12.50	14.29	12.50	10.00	0.00	14.29
PCP				*	30.77	66.67	33.33	10.00	28.57
NDP					*	30.77	35.71	38.46	33.33
TGP						*	50.00	22.22	12.50
AF <sub>1</sub>							*	62.50	37.50
AF <sub>2</sub>								*	25.00
AF <sub>3</sub>									*

Source: Field Survey, 2012






Figure 2: Classification of land use types based on similarity in their key livelihood tree species composition

Source: Field Survey, 2012

## DISCUSSION

Key livelihood tree species richness and alpha diversity were higher in the SNR than any of the introduced land use types. Anthropogenic impacts of habitat destruction have been known to cause biodiversity decay worldwide. Several studies (e.g. Wilson, 1988; Ihuma, *et al.*, 2011; Chima and Omoemu, 2012; Chima and

Uwaegbulem, 2012) lend credence to this assertion. The NDP was next to the SNR in terms of key livelihood tree species richness. About 70% of the tree species found in NDP was among the key livelihood tree species documented by Chima *et al.* (2012). There may be two possible reasons for this. First, NDP has the lowest degree of human-induced

modification, having not been logged since its establishment in 1975. Second, it is located within the residential quarters of the Ogun State Plantation Project in Area J4; the occupants of which may have enhanced the species richness of the key livelihood trees through the dispersal of seeds of eaten fruits. Diversity of the key livelihood tree species was higher in the arable farmlands (especially AF<sub>1</sub> and AF<sub>2</sub>) than in the monoculture plantations except PCP. This could be explained by the high species dominance in the monoculture plantations since diversity takes into account the evenness in the distribution of individuals among the species encountered. It should be noted that *Pinus caribaea* was not listed as one of the key livelihood species, hence diversity was higher and dominance lower in PCP than in other monoculture plantations.

Harris and Silva-Lopez (1992) observed that habitat fragmentation is one of the most serious causes of diminishing biological diversity; while habitat loss is responsible for biodiversity loss and ultimate extinction of species (IUCN, 2002). Thus, the very high ecological distance observed between the SNR and most of the introduced land use types could be attributed to

habitat fragmentation/modification and varying degrees of protection and management. This is made evident in the least similarity recorded between the SNR and CP and the highest between SNR and NDP, when the monoculture plantations were compared with the SNR. Although, the Cocoa plantation is protected, management practices favour only the preferred species while in NDP, diversity of species is tolerated since it acts as a buffer to the residential quarters and not managed for commercial purposes.

In the arable farmlands too (especially AF<sub>1</sub> and AF<sub>2</sub>), more key livelihood tree species were encountered than in most of the monoculture plantations. Apart from the fact that the farms were started as Taungya farms, the farmers also encouraged the growth of trees that contribute to their livelihoods. This explains why AF<sub>1</sub> and AF<sub>2</sub> were more similar to the SNR than most of the monoculture plantations. Also, the closer ecological distance between AF<sub>1</sub> and AF<sub>2</sub> than with AF<sub>3</sub> could be attributed to more years of cultivation in AF<sub>3</sub>. Chima and Omoemu (2012) made a similar observation in tree species composition between a 14-year and 28-year chronosequences of arable farmland, than with

the one that had been under cultivation since over 50-years. However, the closest ecological distance between TGP and PCP, than with any other monoculture plantation, could be attributed to the fact that both sites lie adjacent to each other. The closeness of the sites may have enhanced the exchange of seeds by agents of dispersal.

## CONCLUSION AND

## RECOMMENDATION

This study has shown that natural forest conversion to monoculture plantations and arable farm reduce key livelihood tree species richness and diversity, and that higher degree of

disturbance as a result of high impact logging and longer period of cultivation (beyond thirty years) exacerbates the problem. The absence of *Triplochiton schleroxylon*, *Piptadeniastrum africanum*, *Mansonia altissima*, *Bulchozia coriacea* and *Daniella ogea* (documented as key livelihood tree species) in all land use types enumerated, calls for an all encompassing survey of their populations to include land use/cover types not covered in this study to truly ascertain their level of rarity in the reserve.

## REFERENCES

- Chima, U.D.; Kamalu, O.J. and Omokhua, G.E. (2011). Comparative assessment of floral and faunal diversity of two proposed project sites in the Niger Delta Region of Nigeria. *International Journal of Plant, Animal and Environmental Sciences*, 1(2): 55-62. Available online at [www.ijpaes.com](http://www.ijpaes.com)
- Chima UD, Adedire MO, Aduradola AM, and Agboola DA. (2012). Key livelihood tree species in Omo Biosphere Reserve: A preliminary documentation towards the investigation of land use change impact on key livelihood tree populations. *Greener J Agric Sci* 2(8): 406 – 411.
- Chima, U.D. and Omoemu, F. (2012). Impact of shifting agriculture on tree abundance in an agrarian community within the Niger Delta Region of Nigeria. *ARNP Journal of Agricultural and Biological Science* 7(9): 689 – 698.
- Chima, U.D. and Uwaegbulem, C. 2012. Comparative evaluation of tree species populations under different land use types within the University of Port Harcourt environs. *Tropical Agricultural Research and Extension* 15(2): 1 -7.
- FAO (2006). *Better Forestry, Less Poverty: A practitioner's guide*. FAO Forestry Paper No. 149, FAO, Rome, 66pp.
- Giramet-Carpentier, C., Pelissier, R., Pascal, J. P. and Francois, H. (1998). Sampling strategies for the assessment of tree species diversity. *Journal of Vegetation Science* 9: 161-172.
- Harris, L.D. and Silva-Lopez (1992). Forest fragmentation and the conservation of biological diversity, pp. 197-237. In: P.L. Fielder and S.K. Jain (eds.). *Conservation Biology, The Theory and Practice of Nature Conservation*,

- Preservation and Management*. John Wiley, New York.
- Ihuma, J.O.; Chima, U.D. and Chapman, H.M. (2011). Tree diversity in a Nigerian montane forest ecosystem and adjacent fragmented forests. *ARPJN Journal of Agricultural and Biological Science*, 6(2): 17-22. Available online at [www.arpnjournals.com](http://www.arpnjournals.com).
- Ikhuoria, I.A.; Ero, I.I.; and Ikhuoria, A.E. (2006). Satellite detection and GIS analysis of lowland rainforest reserve reduction in Edo State, Nigeria. In: Salami, A. T. (ed.). *Imperatives of Space Technology for Sustainable Forest Management*. Proceedings of an International Stakeholders' Workshop Sponsored by National Space Research and Development Agency (NARSDA), Abuja.
- Isichei, A. O. (1995). Omo biosphere reserve: current status, utilisation of biological resources and sustainable management. South-South Co-operation on Environmentally Sound Socio-Economic Development in the Humid Tropics, Working Paper 11, UNESCO, Paris, 48pp.
- IUCN (2002). Effective human activities, causes of biodiversity loss (Online). Available by International Union for Conservation of Nature and Natural Resources. <http://www.iucn.org/bil/habitat.html>
- Jansen S. and Vegelius J. (1981). Measurement of ecological association. *Oecologia*. 49: 371-376.
- Keay, R.W.J. (1989). *Trees of Nigeria*, Clarendon Press, Oxford, 475pp.
- Magurran, A. E. (1988). *Ecological Diversity and Its Measurement*. Croom Helm, London.
- Magurran, A.E. (2004). *Measuring Biological Diversity*. Blackwell, Oxford.
- Odum, E.P. (1971). *Fundamentals of Ecology*. W.B Saunders Company, Philadelphia, 574pp.
- Ogunleye, A.J.; Adeola, A.O.; Ojo, L.O. and Aduradola, A.M. (2004). Impact of farming activities on vegetation in Olokemeji Forest reserve, Nigeria, *Global Nest*, 6(2): 131-140.
- Ojo, L.O. (2004). The fate of a tropical rainforest in Nigeria: Abeku sector of Omo Forest Reserve, *Global Nest*, 6(2):116 – 130.
- Ola-Adams, B. A. (1999). Biodiversity Inventory of Omo Biosphere Reserve, Nigeria. Country Report on Biosphere Reserves for Biodiversity Conservation and Sustainable Development in Anglophone Africa (BRAAF) Project. 351 pp.
- Pielou, E.E. (1969). *An Introduction to Mathematical Ecology*. Wiley-Interscience, New York.
- Salami, A. T. (2006). Monitoring Nigerian Forest with NigeriaSat-1 and other satellites pp 28 – 61. In: Salami, A. T. (ed.). *Imperatives of Space Technology for Sustainable Forest Management*. Proceedings of an International Stakeholders' Workshop Sponsored by National Space Research and Development Agency (NARSDA), Abuja.
- Simpson, E.H. (1949). Measurement of diversity, *Nature*, 163:688
- Wilson, E.O. (1988). The current state of biological diversity. pp 3 – 18. In E.O. Wilson, (Ed.). *Biodiversity*. Natural Academy press, Washington, D.C.
- Wolda H. (1983). Diversity, diversity indices and tropical cockroaches. *Oecologia*, 58: 290-298.
- World Bank, (2001). *A revised forest strategy for the World Bank Group*. Draft. World Bank, Washington DC, USA.